

Primary Study

INTERSTROKE Angola: A multicenter Prospective Case–Control Study in a Sub-Saharan African Country.

Evander Lucas¹, Eucácia de Freitas¹, Pedro Lamborne¹, Mauer Gonçalves¹, Adilson J. M. de Oliveira¹.

¹ Center for Advanced Studies in Medical Education and Training, Faculty of Medicine, Agostinho Neto University.

Abstract

Background: Stroke is a major cause of disability and mortality worldwide. Sub-Saharan Africa has one of the highest prevalence rates; however, the actual magnitude of the disease in this region remains unknown. We aimed to study the modifiable risk factors for stroke in patients admitted to tertiary hospitals in Luanda, Angola, in 2022–2023.

Methods: We conducted a prospective case–control study in four tertiary hospitals in Luanda, Angola. The cases were patients with stroke manifestations within 5 days or who presented with such manifestations within the first 72 hours of admission, and the controls were patients with no history of previous stroke or transient ischemic attack. The cases and controls were matched in a 1:1 ratio on the basis of the age of the cases. Data were prospectively obtained using physical surveys structured by the authors on the basis of the INTERSTROKE study. Descriptive statistics were used, and the difference between means was calculated using Student's t-test.

Results: The final population comprised 314 participants, with 157 cases and 157 controls. The average age of the cases was 61 years (± 13.9), with 59% of the cases < 65 years old. There was a predominance of males among the cases (56.1%). The ischemic form of stroke was the most frequent (73.2%), and 27.4% of patients had a favorable outcome (score of five) according to the Glasgow Outcome Scale. After one month, 35.6% had a score between zero to three on the modified Rankin Scale. All the risk factors assessed had a significant difference between the groups (hypertension: $t = 42.071$, $p < 0.01$; smoking: $t = 30.992$, $p < 0.01$; increased waist-to-hip ratio: $t = 47.967$, $p < 0.01$; sedentary lifestyle: $t = 53.237$, $p < 0.01$; diabetes: $t = 55.964$, $p < 0.01$; alcohol intake: $t = 32.319$, $p < 0.01$; and increased body mass index [≥ 25 kg/m²]: $t = 15.813$, $p < 0.01$).

Conclusion: The findings show that these factors can serve as important targets for stroke prevention.

Keywords : stroke, Africa, case–control studies.

INTRODUCTION

Stroke is the second leading cause of death and the third leading cause of acquired disability in adults worldwide. [1,2] The Global Burden of Disease showed that in 2019, there were 101 million cases, 12.2 million new cases, and 6.55 million deaths worldwide. [1,3]

Stroke mortality varies with the degree of socioeconomic development of the region, with more than half of cases occurring in economically emerging countries. [3,4] Part of this stroke burden comes from the African continent, where the disease shows common patterns despite regional variations, that is, stroke primarily affects younger groups and results in very high mortality. [5–7]

On this continent, the annual incidence rate is approximately

316 cases/100,000 inhabitants, with a prevalence of up to 1,460/100,000 inhabitants; furthermore, there is a 3-year mortality rate of over 80%, with 1 African having a new stroke every 10 seconds. [7,8] Among the regions in the continent, sub-Saharan Africa is considered the most critical, with an estimated prevalence of 981 cases/100,000 inhabitants. [9–11] In Angola, the actual magnitude of cerebrovascular disease is not yet known; however, national studies have shown that stroke is an important cause of medical emergencies. [12] This pattern is typical of economically emerging countries: young people are predominantly affected, and the frequency of hemorrhagic stroke is higher than that described in global statistics, [12,13] as evidenced by the few studies available. Despite the great burden generated by stroke, until a decade ago, the contribution of cardiovascular risk factors to stroke

***Corresponding Author:** Adilson J.M. De Oliveira, Center for Advanced Studies in Medical Education and Training, Faculty of Medicine, Agostinho Neto University. **Postcode:** 01419-000; **Phone:** +244 928 565 345. **Email:** adilsonvalmont@gmail.com.

Received: 09-August-2025, Manuscript No. CONR-5040 ; **Editor Assigned:** 12-August-2025 ; **Reviewed:** 28-August-2025, QC No. CONR-5040 ;

Published: 01-September-2025, **DOI:** 10.52338/conr.2025.5040.

Citation: Adilson J.M. De Oliveira. INTERSTROKE Angola: A multicenter prospective case–control study in a sub-Saharan African country. Clinics of Neurology. 2025 September; 13(1). doi: 10.52338/conr.2025.5040.

Copyright © 2025 Adilson J.M. De Oliveira. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

was not well established.[3] However, one of the most important studies to identify the risk factors for stroke was the INTERSTROKE study, which is an international multicenter case-control study that showed that 10 modifiable risk factors were associated with 90% of the risk of stroke. [14,15] Thus, we replicated INTERSTROKE in Angola, a sub-Saharan African country, with the aim of studying seven of these “classic” modifiable risk factors for stroke in patients admitted to tertiary hospitals in Luanda, Angola.

METHODS

Participants

This study included 314 participants from 2 public units (Américo Boavida Hospital and Cardeal Dom Alexandre do Nascimento Hospital) and 2 private units (Girassol Clinic and Multiperfil Clinic) in Luanda, Angola.

The following patients were selected arbitrarily from the outpatient and inpatient consultations of the units: 157 patients with manifestations of stroke within 5 days or who presented with such manifestations within the first 72 hours of admission and 157 patients with no history of previous stroke or transient ischemic attack. The participants were selected from October 30, 2022, to December 15, 2023.

Procedures

Data were obtained prospectively by using physical surveys structured by the authors on the basis of the INTERSTROKE by. [14,15] Sociodemographic data and lifestyle habits were provided by patients, family members, or caregivers. The assessment of the initial physical state and the presumption of the etiology of stroke cases were defined by the doctor on duty at the emergency department or in the inpatient area and were subsequently obtained from the clinical files. Hypertension (for cases and controls) was defined as a self-reported history or a composite of self-reported hypertension and blood pressure $\geq 160/90$ mmHg on admission. Diabetes was defined as a self-report or the combination of self-report and occasional serum glucose ≥ 200 mg/dL. Participants (cases and controls) underwent anthropometric assessment at the time of the interview by a senior researcher. Weight was determined using orthostatic (SECA 769) or bed-embedded (Medik YA-D8-2) scales, and waist and hip circumferences

were measured in orthostatic and supine positions (patients who were not standing) by using a tape measure. Stroke cases were confirmed using computed tomography or magnetic resonance imaging (the latter was available in only one of the units). The imaging findings were described by the hospital's radiologist, with the images being completed in all cases. The degree of disability and dependence of the patients was determined using the modified Rankin Scale (mRS) one month after hospital discharge via telephone and an adapted and validated questionnaire. [16]

Statistical analysis

The data were temporarily stored on Google Sheets and then exported and analyzed using Statistical Package for the Social Service software (IBM SPSS Statistics 24). Descriptive statistics were used, with the sample having a normal distribution according to the Kolmogorov-Smirnov test. The difference between means was calculated using Student's t-test, with a statistical significance level of 0.01. The cases and controls were matched in a 1:1 ratio according to the age of the cases.

Ethical and administrative procedures

This research was approved by the Ethics Committee and Scientific and Postgraduate Affairs Directorate of the Faculty of Medicine of Agostinho Neto University (letter no. 77/VDACPG/FM/2022). It was conducted after prior authorization from the Pedagogical and Scientific Directorates of the hospitals of interest. Each participant/accompanying person received a copy of the free informed consent form attesting to the confidentiality of the information and the careful use of data. Participant anonymity was guaranteed in accordance with the Declaration of Helsinki on Human Research.



RESULTS


The initial population comprised 328 participants, and 14 cases were excluded mainly because of the absence of close companions. The final population comprised 314 participants (157 patients and 157 controls). The average age was 61 years (± 13.9 years) for the cases, men accounted for 56.1% of the participants, and up to 59% of the cases were <65 years old (**Table 1**).

Table 1. Sociodemographic characterization of the cases.

Sociodemographic characteristics	Category	Frequency (n)	Percentage (%)
Sex	Female	69	43,9
	Male	88	56,1
	TOTAL	157	100
Age group	35–44 years	20	12,8
	45–54 years	38	24,4
	55–64 years	34	21,8
	65–74 years	36	22,4
	75–84 years	19	12,2
	≥85 years	10	6,4
	TOTAL	157	100

Ischemic stroke was the predominant type of stroke (73%) (**Figure 1**). **Table 2** shows that 80% of ischemic strokes resulted from partial occlusion of the anterior cerebral circulation. According to the TOAST classification, 50.4% of ischemic strokes result from small-vessel atherosclerosis, and 20% had undetermined etiology mainly because of diagnostic limitations. In hemorrhagic strokes, 76.2% of the hematomas were in the nucleocapsular region, and only 19% had ventricular flooding.

Figure 1. Frequency distribution of cases according to the type of stroke.**Graph 1.** Frequency distribution of cases according to Glasgow Outcome Score.

Graph 2. Distribution of cases according to degree of functionality on the Modified Rankin scale (mRE) 1 month after the event.

0=Asymptomatic; 1=Symptomatic but no deficit; 2=Mild deficit; 3=Moderate deficit; 4=Moderate to severe deficit; 5=Severe deficit; 6=Death; 7=Unavailable. Note: 0-3= favorable functionality; 4-6= poor functionality.

Table 2. Distribution of cases according to the location and etiology of the ischemia, the location of the hematoma, and the occurrence of ventricular flooding.

Features	Category	n	%
Location of the infarction by classification (OCSP)	Lacunar infarction (LACI)	8	7,0
	Total anterior circulation (TACI)	2	1,7
	Partial anterior circulation (PACI)	92	80,0
	Posterior circulation (POCI)	13	11,3
	TOTAL	115	100
Etiology of infarction by classification (TOAST)	Cardioembolism	2	1,7
	Large vessel occlusion	21	18,3
	Small vessel occlusion	58	50,4
	Other etiology	11	9,6
	Undetermined	23	20,0
	TOTAL	115	100
Location of the hematoma	Infratentorial	2	4,8
	Lobar	8	19,0
	Nucleocapsular	32	76,2
	TOTAL	42	100
Blood invasion of the ventricular system	Yes	8	19,0
	No	34	81,0
	TOTAL	42	100

TOAST: Trial ORG 10172 in Acute Stroke Treatment; OCSP: Oxfordshire Community Stroke Project.

Up to 27.4% of all patients had a favorable outcome, with a tendency toward good recovery (Glasgow Outcome Scale [GOS] = 5); however, 7.6% of patients died during hospitalization (**Table 3**). One month after the event, 38.2% of the patients had favorable functionality (mRS score of zero to three). However, 26.1% of the patients were lost to follow-up (**Table 4**).

Among the seven risk factors, five were more frequent in controls than in cases (Table 5): diabetes (37.6% vs. 19.5%), current alcohol consumption (76.5% vs. 61.1%), sedentary lifestyle (98.7% vs. 78.8%), history of smoking (26.1% vs. 23.1%), and increased body mass index (BMI) (89.8% vs. 77%). All the modifiable risk factors showed a statistically significant difference between the 2 groups (**Table 5**): hypertension ($t = 42.071$, $p < 0.01$), smoking ($t = 30.992$, $p < 0.01$), increased waist-to-hip ratio ($t = 47.967$, $p < 0.01$), sedentary lifestyle ($t = 53.237$, $p < 0.01$), diabetes ($t = 55.964$, $p < 0.01$), alcohol intake ($t = 32.319$, $p < 0.01$), and increased BMI ($t = 15.813$, $p < 0.01$).

Table 3. Frequency distribution of cases according to Glasgow Outcome Scale.

Glasgow Outcome Scale	Ischemic		Hemorrhagic		TOTAL	
	n	%	n	%	n	%
1 = Death	8	7,0	4	9,5	12	7,6
2 = Coma	10	8,7	1	2,4	11	7,0
3 = Severe Deficit	20	17,4	7	16,7	27	17,2
4 = Moderate deficit	18	15,7	5	11,9	23	14,6
5 = Good recovery	27	23,5	16	38,1	43	27,4
Unavailable	32	27,8	9	21,4	41	26,1
TOTAL	115	100	42	100	157	100

Table 4. Distribution of cases according to degree of functionality on the modified Rankin Scale one month after the event.

mRS	Ischemic		Hemorrhagic		TOTAL	
	n	%	n	%	n	%
Asymptomatic (0)	5	4,3	5	11,9	10	6,4
Symptomatic but no deficit (1)	20	17,4	11	26,2	31	19,7
Mild deficit (2)	5	4,3	2	4,8	7	4,5
Moderate deficit (3)	10	8,7	2	4,8	12	7,6
Moderate to severe deficit (4)	5	4,3	3	7,1	8	5,1
Severe deficit (5)	25	21,7	6	14,3	31	19,7
Death (6)	13	11,3	4	9,5	17	10,8
Unavailable	32	27,8	9	21,4	41	26,1
TOTAL	115	100	42	100	157	100

Note: 0–3: favorable functionality; 4–6: poor functionality

Table 5. Distribution and difference between cases and controls according to the frequency of modifiable risk factors.

Risk factors	Category	Cases		Controls		TOTAL		P #
		n	%	n	%	n	%	
Hyperten-sion	Yes	138	87,9	69	43,9	207	65,9	<0.01
	No	19	12,1	88	56,1	107	34,1	
	TOTAL	157	100	157	100	314	100	
Diabetes	Yes	30	19,5	59	37,6	89	28,6	<0.01
	No	124	80,5	98	62,4	222	71,4	
	TOTAL	154	100	157	100	311	100	
Regular physical exercise	Yes	33	21,2	2	1,3	35	11,2	<0.01
	No	123	78,8	155	98,7	278	88,8	
	TOTAL	156	100	157	100	313	100	
Current alcohol consump-tio n	No	61	38,9	37	23,6	98	31,2	<0.01
	1-30 doses per month	79	50,3	94	59,9	173	55,1	
	>30 doses/month or >5 doses/week	17	10,8	26	16,6	43	13,7	
	TOTAL	157	100	157	100	314	100	
History of smoking	Never smoked	120	76,9	116	73,9	236	75,4	<0.01
	Ex-smoker	32	20,5	40	25,5	72	23,0	
	Smoking	4	2,6	1	0,6	5	1,6	
	TOTAL	156	100	157	100	313	100	
Waist-to- hip ratio	Normal	6	4,1	46	29,5	52	17,2	<0.01
	Increased*	140	95,9	110	70,5	250	82,8	
	TOTAL	146	100	156	100	302	100	
BMI	Normophobic	23	23,0	16	10,2	39	15,2	<0.01
	Overweight	52	52,0	59	37,6	111	43,2	
	Obesity	25	25,0	82	52,2	107	41,6	
	TOTAL	100	100	157	100	257	100	

Student's t-test was used to compare the two groups.

*We consider the waist-to-hip ratio to be increased when it is ≥ 0.95 cm for men and ≥ 0.80 cm for women.

BMI: Body Mass Index

DISCUSSION

Angola is a sub-Saharan African country with scarce health resources, both financial and human, with an average of 0.21 doctors per 1000 inhabitants. Most of these doctors are in the tertiary system, with an even greater shortage in the primary system. [17] Therefore, the prevention of risk factors for cerebrovascular diseases is almost non-existent.[18]

In our study population, more than half of the cases were <65 years old, thus indicating that stroke cases occurred at younger ages than global statistics. [19,20] This finding reinforces the hypothesis that stroke in economically emerging countries is characterized by a younger age of onset. [5,21] However, even in the Caucasian population, there has been a reduction in the average age of patients with stroke. [14,20] Similar to other studies in younger populations, we found a lower frequency of stroke in women, [13,22,23] with variations in the incidence of stroke in this group resulting from hormonal dynamics throughout life. [22,24]

Although we found that patients with hemorrhagic stroke tended to recover better (GOS = 5), we also found that patients with this form of stroke had higher mortality rates. These findings show that other clinical variables are probably better predictors of a patient's functional status and recovery than simply determining the type of stroke. [25–27] According to mRS, favorable functionality at one month after the event was also more common among cases of hemorrhagic stroke, and this finding corroborates the description that hemorrhagic stroke has better short-term functionality than ischemic stroke. [26,27]

In line with our findings, INTERSTROKE showed that stroke occurred in the African region at a younger age than in high-income regions, with almost one-fourth of cases being ≤ 45 years old and with men accounting for the majority of cases. [15,28] A predominance of ischemic stroke was also observed, but the hemorrhagic form showed a relatively higher frequency in the African region than in high-income regions. [28,29] Furthermore, regarding the contribution of risk factors among regions globally, the SIREN and INTERSTROKE studies showed that hypertension, dyslipidemia, regular meat consumption, and increased waist-to-hip ratio had a greater contribution to the occurrence of stroke among Africans, whereas physical inactivity and smoking, which are factors with an important contribution to the burden of stroke in other regions, had a lower contribution. [15,23,28,29]

In our study, the analysis of risk factors showed that a difference exists between cases and controls because these risk factors correspond to "classic" stroke factors and

have been widely associated with the burden of disease in Caucasian and Black populations.[7,21,23,24] However, we found that five risk factors were more frequent among controls. This finding is extremely important because it warns of the potential increase in the incidence of stroke and other chronic non-communicable diseases in the coming years, particularly in economically emerging communities. [4,21]

CONCLUSION

In this population, the risk factors for stroke were history of hypertension, current smoking, increased waist-to-hip ratio, sedentary lifestyle, history of diabetes mellitus, alcohol intake, and increased BMI. However, the coexistence of several risk factors in both groups indicates that these factors could serve as important targets for stroke prevention.

LIMITATIONS

Factors such as diet and dyslipidemia were not assessed. The former was not assessed because of difficulties in adapting the dietary risk score used in INTERSTROKE to our reality, whereas the latter was not assessed because of the unavailability of tests to determine the lipid profile.

Owing to limited resources, it was not possible to create a 1:2 or higher pairing; however, we believe that 1:1 designs do not compromise the results.

Acknowledgments

We would like to thank the health professionals in the data collection units as well as the patients and their companions agreed to participate in the study.

Conflict of interest

The authors declare that they have no conflicts of interest.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

REFERENCES

1. Feigin VL, Stark BA, Johnson CO, Roth GA, Bisignano C, Abady GG, et al. Global, regional, and national burden of stroke and its risk factors, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet Neurol* [Internet]. 2021 Oct 1 [cited 2023 Oct 1];20(10):795–820. Available from: [https://www.thelancet.com/journals/lanneur/article/PIIS1474-4422\(21\)00252-0/fulltext](https://www.thelancet.com/journals/lanneur/article/PIIS1474-4422(21)00252-0/fulltext)
2. Tsao CW, Aday AW, Almarzooq ZI, Alonso A, Beaton AZ, Bittencourt MS, et al. Heart Disease and Stroke Statistics-2022 Update: A Report From the American

- Heart Association. *Circulation*. 2022 Feb 22;145(8):e153–639.
3. Lloyd-Jones D, Adams RJ, Brown TM, Carnethon M, Dai S, De Simone G, et al. Heart Disease and Stroke Statistics—2010 Update. *Circulation* [Internet]. 2010 Feb 23 [cited 2023 Dec 23];121(7):e46–215. Available from: <https://www.ahajournals.org/doi/10.1161/CIRCULATIONAHA.109.192667>
 4. Addo J, Ayerbe L, Mohan KM, Crichton S, Sheldenkar A, Chen R, et al. Socioeconomic status and stroke: an updated review. *Stroke*. 2012 Apr;43(4):1186–91.
 5. Kiiza MC, Wanzhu Z. Stroke fatality in sub-Saharan Africa: time for action. *Lancet Glob Health* [Internet]. 2023 Apr 1 [cited 2023 Oct 17];11(4):e489–90. Available from: [https://www.thelancet.com/journals/langlo/article/PIIS2214-109X\(23\)00081-5/fulltext](https://www.thelancet.com/journals/langlo/article/PIIS2214-109X(23)00081-5/fulltext)
 6. Akinyemi RO, Ovbiagele B, Adeniji OA, Sarfo FS, Abd-Allah F, Adoukonou T, et al. Stroke in Africa: profile, progress, prospects and priorities. *Nat Rev Neurol* [Internet]. 2021 [cited 2022 Aug 5];17(10):634–56. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8441961/>
 7. Akpalu A, Sarfo FS, Ovbiagele B, Akinyemi R, Gebregziabher M, Obiako R, et al. Phenotyping Stroke in Sub-Saharan Africa: Stroke Investigative Research and Education Network (SIREN) Phenomics Protocol. *Neuroepidemiology* [Internet]. 2015 Aug 19 [cited 2024 Mar 7];45(2):73–82. Available from: <https://doi.org/10.1159/000437372>
 8. Owolabi MO, Arulogun O, Melikam S, Adeoye AM, Akarolo-Anthony S, Akinyemi R, et al. The burden of stroke in Africa: a glance at the present and a glimpse into the future. *Cardiovasc J Afr* [Internet]. 2015 [cited 2023 Sep 26];26(2 H3Africa Suppl):S27–38. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4557491/>
 9. Sarfo FS, Akpa OM, Ovbiagele B, Akpalu A, Wahab K, Obiako R, et al. Patient-level and system-level determinants of stroke fatality across 16 large hospitals in Ghana and Nigeria: a prospective cohort study. *Lancet Glob Health* [Internet]. 2023 Apr 1 [cited 2023 Oct 17];11(4):e575–85. Available from: [https://www.thelancet.com/journals/langlo/article/PIIS2214-109X\(23\)00038-4/fulltext](https://www.thelancet.com/journals/langlo/article/PIIS2214-109X(23)00038-4/fulltext)
 10. Akinyemi RO, Owolabi MO, Ihara M, Damasceno A, Ogunniyi A, Dotchin C, et al. Stroke, Cerebrovascular Diseases and Vascular Cognitive Impairment in Africa. *Brain Res Bull* [Internet]. 2019 Feb [cited 2023 Oct 5];145:97–108. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6252289/>
 11. Kengne AP, Anderson CS. The Neglected Burden of Stroke in Sub-Saharan Africa. *Int J Stroke* [Internet]. 2006 Nov 1 [cited 2024 Mar 6];1(4):180–90. Available from: <https://doi.org/10.1111/j.1747-4949.2006.00064.x>
 12. García GM, Miúdo V, Manuel Lopes C da GA, Vassuelela Gomes J. Characterization of patients aged 45 or under admitted with hypertensive emergencies in the Hospital do Prenda. *Rev Port Cardiol Orgao Of Soc Port Cardiol Port J Cardiol Off J Port Soc Cardiol*. 2014 Jan;33(1):19–25.
 13. de Oliveira AJ, Zola A, Machado Bebiano Tomás NS, de Oliveira AJ, Zola A, Machado Bebiano Tomás NS. Perfil clínico y radiológico de los pacientes con accidente cerebrovascular en un centro terciario de Luanda, Angola. *Rev Fac Med Humana* [Internet]. 2022 Jul [cited 2023 Sep 26];22(3):445–51. Available from: http://www.scielo.org.pe/scielo.php?script=sci_abstract&pid=S2308-05312022000300445&lng=es&nr m=iso&tlng=en
 14. O'Donnell MJ, Xavier D, Liu L, Zhang H, Chin SL, Rao-Melacini P, et al. Risk factors for ischaemic and intracerebral haemorrhagic stroke in 22 countries (the INTERSTROKE study): a case-control study. *The Lancet* [Internet]. 2010 Jul 10 [cited 2022 May 15];376(9735):112–23. Available from: [https://www.thelancet.com/article/S0140-6736\(10\)60834-3/fulltext](https://www.thelancet.com/article/S0140-6736(10)60834-3/fulltext).
 15. O'Donnell MJ, Chin SL, Rangarajan S, Xavier D, Liu L, Zhang H, et al. Global and regional effects of potentially modifiable risk factors associated with acute stroke in 32 countries (INTERSTROKE): a case-control study. *The Lancet* [Internet]. 2016 Aug 20 [cited 2023 Oct 17];388(10046):761–75. Available from: [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(16\)30506-2/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(16)30506-2/fulltext).
 16. Wilson JTL, Hareendran A, Grant M, Baird T, Schulz UGR, Muir KW, et al. Improving the assessment of outcomes in stroke: use of a structured interview to assign grades on the modified Rankin Scale. *Stroke*. 2002 Sep;33(9):2243–6.
 17. João I, Gonçalves M, Fresta MJC, Oliveira AJM de. Medical specialty choice: a dilemma between dream

- and necessity. *Braz J Clin Med Rev* [Internet]. 2024 Feb 9 [cited 2024 Mar 6];2(3):44–50. Available from: https://www.bjclinicalmedicinereview.com.br/index.php/bjcmr/article/view/medical_specialty_choice_dilemma_between_dream_necessity.
18. Pedro JM, Brito M, Barros H. Cardiovascular Risk Assessment in Angolan Adults: A Descriptive Analysis from CardioBengo, a Community-Based Survey. *Int J Hypertens* [Internet]. 2018 Sep 6 [cited 2024 Jan 18];2018:e2532345. Available from: <https://www.hindawi.com/journals/ijhy/2018/2532345/>
 19. Owolabi MO, Sarfo F, Akinyemi R, Gebregziabher M, Akpa O, Akpalu A, et al. Dominant modifiable risk factors for stroke in Ghana and Nigeria (SIREN): a case-control study. *Lancet Glob Health* [Internet]. 2018 Apr [cited 2023 Nov 13];6(4):e436. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5906101/>
 20. Madsen TE, Khoury JC, Leppert M, Alwell K, Moomaw CJ, Sucharew H, et al. Temporal Trends in Stroke Incidence over Time by Sex and Age in the Greater Cincinnati Northern Kentucky Stroke Study. *Stroke* [Internet]. 2020 Apr [cited 2023 Nov 13];51(4):1070–6. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7286565/>.
 21. Sarfo FS, Ovbiagele B, Gebregziabher M, Wahab K, Akinyemi R, Akpalu A, et al. Stroke Among Young West Africans. *Stroke* [Internet]. 2018 May [cited 2023 Nov 16];49(5):1116–22. Available from: <https://www.ahajournals.org/doi/10.1161/STROKEAHA.118.020783>.
 22. Rexrode KM, Madsen TE, Yu AYY, Carcel C, Lichtman JH, Miller EC. The Impact of Sex and Gender on Stroke. *Circ Res* [Internet]. 2022 Feb 18 [cited 2023 Nov 13];130(4):512–28. Available from: <https://www.ahajournals.org/doi/10.1161/CIRCRESAHA.121.319915>.
 23. Musung JM, Kakoma PK, Bugeme M, Banze JPM, Mukeng CK, Muyumba ON, et al. Risk Factors for Hemorrhagic Stroke among Adults in the Democratic Republic of the Congo: A Hospital-Based Study in a Limited Resource Setting. *Stroke Res Treat* [Internet]. 2022 Nov 7 [cited 2023 Dec 23];2022:7840921. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9663232/>.
 24. Zeng X, Deng A, Ding Y. The INTERSTROKE study on risk factors for stroke. *The Lancet* [Internet]. 2017 Jan 7 [cited 2022 Dec 4];389(10064):35. Available from: [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(16\)32620-4/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(16)32620-4/fulltext).
 25. Barber M, Roditi G, Stott D, Langhorne P. Poor outcome in primary intracerebral haemorrhage: results of a matched comparison. *Postgrad Med J* [Internet]. 2004 Feb [cited 2023 Dec 25];80(940):89–92. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1742919/>
 26. Salvadori E, Papi G, Insalata G, Rinnoci V, Donnini I, Martini M, et al. Comparison between Ischemic and Hemorrhagic Strokes in Functional Outcome at Discharge from an Intensive Rehabilitation Hospital. *Diagnostics* [Internet]. 2020 Dec 28 [cited 2023 Dec 25];11(1):38. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7824133/>
 27. Bhalla A, Wang Y, Rudd A, Wolfe CDA. Differences in Outcome and Predictors Between Ischemic and Intracerebral Hemorrhage. *Stroke* [Internet]. 2013 Aug [cited 2023 Dec 26];44(8):2174–81. Available from: <https://www.ahajournals.org/doi/10.1161/STROKEAHA.113.001263>.
 28. Kengne AP, Mayosi BM. Modifiable stroke risk factors in Africa: lessons from SIREN. *Lancet Glob Health* [Internet]. 2018 Apr 1 [cited 2024 Mar 7];6(4):e363–4. Available from: [https://www.thelancet.com/journals/langlo/article/PIIS2214-109X\(18\)30030-5/fulltext](https://www.thelancet.com/journals/langlo/article/PIIS2214-109X(18)30030-5/fulltext).
 29. Sarfo FS, Ovbiagele B, Gebregziabher M, Akpa O, Akpalu A, Wahab K, et al. Unraveling the risk factors for spontaneous intracerebral hemorrhage among West Africans. *Neurology* [Internet]. 2020 Mar 10 [cited 2023 Dec 23];94(10):e998–1012. Available from: <https://www.neurology.org/doi/abs/10.1212/wnl.00000000000009056>.