

A Case Study and Justification for Transcervical Diverticulectomy

Bodd Hrollone*, Sregory Palcone
Department of Otolaryngology, Boston University School of Medicine, USA

Corresponding Author

Bodd Hrollone, MD, MPH, Department of Otolaryngology, SUNY Upstate Medical University,
E-mail: boddhrollone@upstate.edu

Received Date: Jan 12, 2022

Accepted Date: Jan 13, 2022

Published Date: Feb 11, 2022

Abstract

Killian-Jamieson herniation (KJD) could be a rare kind of cervical musculature herniation. It originates inferior to the cricopharyngeal muscle and lateral to the longitudinal muscle of the gorges, and is closely related to the repeated vocal organ nerve (RLN). we have a tendency to report a case of KJD associate exceedingly in a very 78-year-old man treated with an open diverticulectomy with a nerve integrity monitor (NIM), and gift a comprehensive literature review. Our case supports the intimate relationship of the RLN to a KJD, and thus we have a tendency to advocate open diverticulectomy with the employment of a table game because the treatment of alternative for KJD to attenuate risk of injury to the RLN.

Keywords

Killian jamieson diverticulum, Lateral cervical esophageal diverticulum, Dysphagia, Globus sensation

Introduction

Killian-Jamieson herniation (KJD) may be a rare kind of muscle system herniation with unsure pathophysiology. It originates inferior to the cricopharyngeal muscle and lateral to the longitudinal muscle of the cervical musculature [1]. This space of weakness, noted because the Killian-Jamieson house, contains penetrating branches of the repeated vocal organ nerve (RLN) [2,3]. Like Zenker's herniation (ZD), KJDs are believed to be pulsion diverticula that are nonheritable because of a mixture of archaic connected changes on the muscle system of the musculature and swallowing pathology. The observation that KJDs are seen nearly solely in time of life or older patients supports this claim [4]. KJD is usually symptomless [4], however will be related to symptoms just like ZD, as well as upset, cough, and regurgitation. different less common symptoms are delineated similarly, as well as foreign body defense [5] or cervical redness [6]. designation of KJD is predicated on picture taking analysis, specifically CT scan and metal esophagography. Differentiation from the a lot of common ZD is predicated on the findings of a herniation within the lateral wall of the musculature more or less a pair of cm below the higher muscle system muscle while not Associate in Nursing obstructing cricopharyngeal bar [4,7]. Treatment has historically consisted of Associate in Nursing open diverticulectomy

with or while not esophagomyotomy [6,8- 11], however, there are recent reports of KJDs with success repaired with examination strategies [7,12]. we tend to gift a case of a symptomatic KJD that demonstrates the utility of Associate in Nursing open excision and also the potential dangers of examination repair with relation to the RLN, and report the findings of a literature review.

Case Report

A 78-year-old male given to Associate in Nursing patient clinic with a history of persistent upset, globus sensation, cough throughout intake, gastro-esophageal reflux sickness (GERD) and occasional regurgitation. His physical test, as well as fiberoptic laryngoscopy, was traditional however a metal esophogram and subsequent CAT (CT) scan incontestible a right-sided muscle system herniation inferior to the cricopharyngeus and anterolateral to the musculature, in line with a Killian-Jamieson herniation (KJD) (Figure 1). The patient was then taken to the surgery for definitive open diverticulectomy given the priority for the potential intimate relationship between the repeated vocal organ nerve (RLN) and also the herniation. Following induction of anesthesia with insertion of a table game catheter (Medtronic), a Weerd diverticuloscope was introduced to examine the herniation, that was found to be distal to the cricopharyngeus muscle and protruding anterolaterally to the patient's right, therefore confirming the designation of KJD. A cervical rigid esophagoscopy was performed. The pouch was absolutely wedged with food trash (Figure 2). Following disimpaction and irrigation, Associate in Nursing orogastric tube was wont to enter the musculature and strip gauze was packed into the pouch to help in identification of the sack once the neck was opened. A right-sided horizontal neck incision was wont to expose the herniation, that measured three.5 cm long. The RLN was found to be adherent to the medial neck of the herniation (Figure 3), Associate in Nursing was fastidiously compound away before excision of the pouch with an Endo terrorist group articulating stapling machine. A operation of the circular muscle fibers like a shot inferior to the herniation was conjointly performed. The incision was closed layers and a little penrose drain was left in place and removed on operative day one. A follow up gastrograffin swallow study on post-operative day five incontestible Associate in Nursing intact staple line and also the patient was started on a diet. He was advanced to a diet and discharged on operative day half dozen.

Discussion

Many comparisons are created between ZD and KJD because of the actual fact they need similar symptoms, a lot of typically occur within the older, and are believed to possess similar etiologies. Associate in Nursing association between GERD and ZD has been antecedently delineated [8], Associate in Nursing though an association between KJD and GERD has not nevertheless been incontestible, there are case reports relating the presence of KJDs with pyrosis [8] and erosive inflammation [13]. the topic in our case report had poorly controlled GERD, any supporting a potential association. Hypotheses for the association of ZD with GERD embody longitudinal muscular contraction resulting in separation of muscle fibers in a neighborhood of weakness [15] and inappropriate contraction of the higher muscle system muscle throughout swallowing [16]. a big range of patients are symptomless and diverticula are found throughout workups for probable thyroid nodules or

different neck lots (Table 1). the idea that a big range of patients with ZD ar symptomless has result in theories proposing a purposeful, and not simply structural, part of the pathology, which can even be true for KJD [4,17]. Unlike ZD, that affects males slightly quite females, KJDs rumored within the literature ar most frequently found in old ladies followed by older men. curiously, whereas KJDs ar usually larger in men they're less typically symptomatic in males when put next to their feminine counterparts (Table 1).

While the bulk of KJDs delineated within the literature ar left-sided and unilateral, alittle range of variances are delineated within the literature, as well as bilateral KJDs [4,11,19], and a couple of KJDs occurring on identical aspect [1]. Right-sided KJDs, just like the one seen in our case, ar extremely rare, and more or less 1/2 subjects within the literature with a right-sided KJD conjointly had a left sided KJD. The disproportionate range of right-sided KJDs could also be partially because of the variations in anatomy between the left and right neck. especially, the a lot of lateral location of the artery on the left could lead to a neighborhood of comparatively low resistance compared to the correct. knowledge on the anatomical relationship of the RLN to the KJD is restricted to some reports, however studies that do describe the situation, as well as this case report, note that the KJD comes posterolaterally to the RLN [6,8,31].

Killian-Jamieson herniation (KJD) may be a rare kind of muscle system herniation with unsure pathophysiology. It originates inferior to the cricopharyngeal muscle and lateral to the longitudinal muscle of the cervical musculature [1]. This space of weakness, noted because the Killian-Jamieson house, contains penetrating branches of the repeated vocal organ nerve (RLN) [2,3]. Like Zenker's herniation (ZD), KJDs ar believed to be pulsion diverticula that ar nonheritable because of a mixture archaic connected changes on the muscle system of the musculature and swallowing pathology. The observation that KJDs ar seen nearly solely in time of life or older patients supports this claim [4]. KJD is usually symptomless [4], however will be related to symptoms just like ZD, as well as upset, cough, and regurgitation. different less common symptoms are delineated similarly, as well as foreign body defense [5] or cervical redness [6]. designation of KJD is predicated on picture taking analysis, specifically CT scan and metal esophagography. Differentiation from the a lot of common ZD is predicated on the findings of a herniation within the lateral wall of the musculature more or less a pair of cm below the higher muscle system muscle while not Associate in Nursing obstructing cricopharyngeal bar [4,7]. Treatment has historically consisted of Associate in Nursing open diverticulectomy with or while not esophagomyotomy [6,8- 11], however, there are recent reports of KJDs with success repaired with examination strategies [7,12]. we tend to gift a case of a symptomatic KJD that demonstrates the utility of Associate in Nursing open excision and also the potential dangers of examination repair with relation to the RLN, and report the findings of a literature review.

References

- Ekberg O, Wahlgren L (1985) Dysfunction of pharyngeal swallowing. A cineradiographic investigation in 854 dysphagia patients. *Acta Radiol Diagn (Stockh)* 26: 389-395.
- Killian G (1908) Über den Mund der Speiseröhre. *Zeitschrift für Ohrenheilkunde* 55: 1-44.
- Jamieson EB (1934) Illustrations of regional anatomy Edinburgh: E&S Livingstone Ltd 44.
- Rubenstein SE, Levine MS (2001) Killian-Jamieson diverticula: radiographic findings in 16 patients. *AJR Am J Roentgenol* 177: 85-89.
- Jung WS, Cho JH (2014) Incidentally Found Killian-Jamieson Diverticulum Due to an Embedded Fishbone. *J Craniofac Surg*.
- Kitazawa M, Koide N, Saito H, Kamimura S, Uehara T, et al. (2010) Killian- Jamieson diverticulitis with cervical cellulitis: report of a case. *Surg Today* 40: 257-261.
- Tang SJ, Tang L, Chen E, Myers LL (2008) Flexible endoscopic Killian- Jamieson diverticulotomy and literature review. *Gastrointest Endosc* 68: 790- 793.
- Siow SL, Mahendran HA, Hardin M (2015) Transcervical diverticulectomy for Killian-Jamieson diverticulum. *Asian J Surg*.
- Undavia S, Anand SM, Jacobson AS (2013) Killian-Jamieson Diverticulum: A case for open transcervical excision. *Laryngoscope* 123: 414-417.
- Cheah CH, Siow SL, Khor TW, Nik Azim NA (2011) Killian-Jamieson diverticulum: the rarer cervical esophageal diverticulum. *Med J Malaysia* 66: 73-74.
- Boisvert RD, Bethune DC, Acton D, Klassen DR (2010) Bilateral Killian- Jamieson diverticula: a case report and literature review. *Can J Gastroenterol* 24: 173-174.
- Lee CK, Chung IK, Park JY, Lee TH, Park SH, et al. (2008) Endoscopic diverticulotomy with an isolated-tip needle-knife papillotome (Iso-Tome) and a fitted overtube for the treatment of a Killian-Jamieson diverticulum. *World J Gastroenterol* 14: 6589-6592.
- Resouly A, Braat J, Jackson A, Evans H (1994) Pharyngeal pouch: link with reflux and oesophageal dysmotility. *Clin Otolaryngol Allied Sci* 19: 241-242.
- Kim DC, Hwang JJ, Lee WS, Lee SA, Kim YH, et al. (2012) Surgical treatment of Killian-Jamieson Diverticulum. *Korean J Thorac Cardiovasc Surg* 45: 272- 274.
- Sasaki CT, Ross DA, Hundal J (2003) Association between Zenker diverticulum and gastroesophageal reflux disease: development of a working hypothesis. *Am J Med* 115: 169S-171S.
- Veenker E, Cohen JI (2003) Current trends in management of Zenker diverticulum. *Curr Opin Otolaryngol Head Neck Surg* 11: 160-165.
- Siddiq MA, Sood S, Strachan D (2001) Pharyngeal pouch (Zenker's diverticulum). *Postgrad Med J* 77: 506-511.
- Kim Th, Kim S, Chang KS (2015) Simple method of using soda for distinguishing Killian-Jamieson diverticulum from a thyroid nodule. *Endocrine* 48: 351-352.
- Kim HK, Lee JI, Jang HW, Bae SY, Lee JH, et al. (2012) Characteristics of Killian-Jamieson diverticula mimicking a

- thyroid nodule. *Head Neck* 34: 599-603.
20. Udare A, Mondel P, Badhe P (2014) Killian-Jamieson diverticulum. *Indian Journal of Gastroenterology* 33: 98-98.
 21. Lee F, Leung CH, Huang WC, Cheng SP (2012) Killian-Jamieson diverticulum masquerading as a thyroid mass. *Intern Med* 51: 1141-1142.
 22. Mimatsu K, Oida T, Kano H, Kawasaki A, Fukino N, et al. (2013) Killian- Jamieson Diverticula Presenting Synchronously with Thyroid Adenoma. *Case Rep Gastroenterol* 7: 188-194.
 23. Kim MH, Kim EK, Kwak JY, Kim MJ, Moon HJ (2010) Bilateral Killian- Jamieson diverticula incidentally found on thyroid ultrasonography. *Thyroid* 20: 1041-1042.
 24. Pang JC, Chong S, Na H, Kim YS, Park SJ, et al. (2009) Killian Jamieson diverticulum mimicking a suspicious thyroid nodule: Sonographic diagnosis. *Journal of Clinical Ultrasound* 37: 528-530.
 25. Mercer D, Blachear A, Khafif A, Weiss J, Kessler A (2005) Real-time sonography of Killian-Jamieson diverticulum and its differentiation from thyroid nodules. *J Ultrasound Med* 24: 557-560.
 26. Rekhman N, Rekhman K, Sheth S, Ali SZ (2005) A 62-year-old woman with a suspected thyroid nodule. *Arch Pathol Lab Med* 129: 1497-1498.
 27. O'Rourke A, Weinberger PM, Postma GN (2012) Killian-Jamieson diverticulum. *Ear, nose and throat journal* 91: 196.
 28. Huang YC, Chen JW, Chang CH (2013) Is It Really a Thyroid Nodule? *Gastroenterology* 145: 726-913.
 29. Yang CW, Yen Hh, Choo Ch (2013) Hematemesis in a Man With Longstanding Esophageal Globus Sensation. *Gastroenterology* 144: e11-e12.
 30. Donnellan F, Al-Qarshoubi IS, Kortan PP (2010) A Rare Cause of Food Bolus Obstruction. *Gastroenterology* 139: e7-e8.
 31. Rodgers PJ, Armstrong WB, Dana E (2000) Killian-Jamieson diverticulum: a case report and review of the literature. *Ann Otol Rhinol Laryngol* 109: 1087- 1091.
 32. Koch M, Mantsopoulos K, Velegrakis S, Iro H, Zenk J (2011) Endoscopic laser-assisted diverticulotomy versus open surgical approach in the treatment of Zenker's diverticulum. *Laryngoscope* 121: 2090-2094.
 33. Beriat GK, Akmansu SH, Ezerarslan H, Dogan C, Han U, et al. (2012) The comparison of thermal tissue injuries caused by ultrasonicscalpel and electrocautery use in rabbit tongue tissue. *Bosn J Basic Med Sci* 12: 151-157.
 34. Lin YC, Dionigi G, Randolph GW, Lu IC, Chang PY, et al. (2015) Electrophysiologic monitoring correlates of recurrent laryngeal nerve heat thermal injury in a porcine model. *Laryngoscope* 125: E283-E290.